

Biostimulants, activators, foliar fertilizers.

AMINOPRIM 1.5 L/ha – (regeneration activator) used after stress conditions occur, and also in combination with fungicides or insecticides (after checking compatibility for mixing)

Microbiological products

Pre-plant soil spraying

BACTIM
NEMABLOCK SL
1.2 L/ha

PLONVIT NITRO
2-4 kg/ha
AMINO ULTRA Cu-24
0.3 kg/ha
AMINO ULTRA Mn-22
0.3 kg/ha
MOLYBEDENUM
1 L/ha
VANADOO
0.25 L/ha
TYTANIT
0.2 L/ha

13 — 15

PLONVIT OPTY
2-4 kg/ha
MIKROKOMPLEX
5 kg/ha
AMINO ULTRA Mn-22
0.3 kg/ha

21 — 29

The fertilizer program requires the application of **PROAQUA** at 1 L/ha
PLONVIT OPTY
2 kg/ha
BORON TURBO
0.8 L/ha
AMINO ULTRA Cu-24
0.2 kg/ha
AMINO ULTRA Fe-20
0.2 kg/ha
MOLYBEDENUM
0.5 L/ha
VANADOO
0.25 L/ha
TYTANIT
0.2 L/ha


PLONVIT PHOSPHO
2-4 kg/ha
BORON TURBO
0.8 L/ha
OPTYSIL
0.5 L/ha

51 — 55

PLONVIT KALI
2-4 kg/ha

71 — 73

Recommended medium-droplet spraying
After spraying – crop residues shallowly mix
with the soil



The ideal conditions for foliar spraying are an air temperature of 12–20°C, relative humidity of 60–80%, and a spray solution temperature of 15–20°C. Spraying should be carried out in the early morning or evening. It is essential to strictly adhere to the recommended application rate and concentration for each product individually, taking into account the prevailing weather conditions.