


Seedling production

Biostimulants, activators, foliar fertilizers.


Recommended doses of INTERMAG products for seedling production:	
Product	Application method / Solution concentration
ROOTSTAR	Fertigation / Foliar application: 0.05–0.1%
PLONVIT NITRO PLONVIT OPTY GROWON, BORON TURBO	Fertigation / Foliar application: 0.05–0.1%
AMINO ULTRA Fe-20	Foliar application: 0.05%
TYTANIT	Fertigation / Foliar application: 0.02–0.03%
AMINOPRIM	Foliar application: 0.3%
OPTYSIL	Fertigation / Foliar application: 0.05%



Open-field cultivation


AMINOPRIM 1.5 L/ha (0.2–0.5 1/100 water) - apply after stress conditions occur, as well as in combination with fungicides or insecticides (after checking mixing compatibility).

PROAQUA water conditioner enhances water quality by improving its physicochemical properties and ensuring greater solution stability



Microbiological products

Pre-plant soil spraying
BACTIM
NEMABLOCK SL
1.2 L/ha



The ideal conditions for foliar spraying are an air temperature of 12–20°C, relative humidity of 60–80%, and a spray solution temperature of 15–20°C. Spraying should be carried out in the early morning or evening. It is essential to strictly adhere to the recommended application rate and concentration for each product individually, taking into account the prevailing weather conditions.