

AMINOPRIM 1.5 L/ha – (regeneration activator) used after stress conditions occur, and also in combination with fungicides or insecticides (after checking compatibility for mixing)

SULVIT
5 L/ha

ROOTSTAR
1.2 L/ha

PLONVIT PHOSPHO
4 kg/ha
AMINO ULTRA HORTI
0.75 kg/ha
TYTANIT
0.2 L/ha
VANADOO
0.25 L/ha

PLONVIT NITRO
4 kg/ha
SULVIT
5 L/ha
AMINO ULTRA Zn-24
0.55 kg/ha
AMINO ULTRA Mn-22
0.4 L/ha
OPTYSIL
0.5 L/ha

PLONVIT KALI
4 kg/ha
AMINO ULTRA HORTI
0.75 kg/ha
AMINO ULTRA Cu-24
0.6 kg/ha
MOLYBDEDENUM
1 L/ha
TYTANIT
0.2 L/ha
OPTYSIL
0.5 L/ha

KALPIM
5 L/ha
AMINO ULTRA HORTI
0.75 kg/ha
TYTANIT
0.2 L/ha
VANADOO
0.25 L/ha

KALPIM
5 L/ha
AMINO ULTRA Cu-24
0.6 kg/ha
OPTYSIL LONGLIFE
1.5 kg/ha

Pre-plant soil spraying
BACTIM
NEMABLOCK SL
1.2 L/ha

Soil preparation for
planting garlic cloves

Before planting garlic
cloves

Planting garlic cloves

Winter dormancy

Spring: After plant
emergence

leaf stage 5–3

Leaf growth

Beginning of garlic bulb
formation

Garlic bulb reaches
30–50% of typical size

Garlic bulb reaches
60–80% of typical size

Harvest

The ideal conditions for foliar spraying are an air temperature of 12–20°C, relative humidity of 60–80%, and a spray solution temperature of 15–20°C. Spraying should be carried out in the early morning or evening. It is essential to strictly adhere to the recommended application rate and concentration for each product individually, taking into account the prevailing weather conditions.